

| Cat.Nr. | Product  | Rxn per kit |
|---------|--|-------------|
|         | HIV  |             |
|         | HIV-1 p24 – antigen – EIA - BEST   |             |
| D-0134  | Enzyme immunoassay kit for the detection and confirmation of HIV-1 p24 antigen presence.                                     | 12x8        |
| D-0151  | CombiBest HIV - 1,2 AG/AB (set 1)  | 24x8        |
| D-0152  | CombiBest HIV - 1,2 AG/AB (set 2)  | 12x8        |
|         | Enzyme immunoassay kit for the detection of anti-HIV - 1,2 antibodies and HIV-1 p24 antigen.                                 |             |
| D-0193  | UniBest HIV – 1,2 Ab (set 1)   | 24x8        |
| D-0192  | UniBest HIV - 1,2 Ab (set 2)   | 12x8        |
|         | Enzyme immunoassay kit for the detection of total antibodies against HIV-1,2.  |             |
|         | LIA – blot - HIV - 1,2 BEST  |             |
| D-0160  | Reagent kit for the detection of antibodies against HIV-<br>1,2 antigens by immunoblot.                                      | 24          |
|         | HEPATITIS A  |             |
|         | Vectohep A - IgM   |             |
| D-0352  | Enzyme immunoassay kit for the detection of IgM against hepatitis A virus.   | 12x8        |
|         | HAV – antigen – EIA - BEST   |             |
| D-0356  | Enzyme immunoassay kit for the detection of hepatitis A virus antigen.   | 12x8        |
| D-0362  | Vectohep A - IgG<br>Enzyme immunoassay kit for the detection of IgG against<br>hepatitis A virus.                            | 12x8        |
|         | HEPATITIS B  |             |
| D-0543  | HBsAg – EIA – BEST (set 2)   | 192         |
| D-0544  | HBsAg – EIA – BEST (set 3)   | 12x8        |
|         | Enzyme immunoassay kit for the detection of HBs-<br>antigen. Sensitivity – 0.01 ng/ml.                                       |             |
|         | HBsAg – EIA – BEST (quantitative analysis)   |             |
| D-0545  | Enzyme immunoassay kit for the quantitative analysis of HBs-antigen. Sensitivity – 0.05 ng/ml. Measurement range: 0-10ng/ml. | 12 x 8      |

## RUO ELISA reagent list for 2016 (not for CIS countries)

|        | HBsAg – confirmatory – EIA - BEST   |      |
|--------|---|------|
| D-0546 | Enzyme immunoassay kit for the confirmation of HBs-<br>antigen presence. Sensitivity – 0.01 ng/ml.  | 6x8  |
| D-0557 | Vectohep B – HBs – antigen (set 1)  | 24x8 |
| D-0556 | Vectohep B – HBs – antigen (set 3)  | 12x8 |
|        | Enzyme immunoassay kit for the detection of HBs-<br>antigen. Sensitivity – 0.05 ng/ml.  |      |
| D-0558 | Vectohep B-HBs – antigen (set 5)<br>Enzyme immunoassay kit for the detection and<br>confirmation of HBs-antigen presence. Sensitivity – 0.05<br>mg/ml.  | 6x8  |
| D-0562 | VectoHBsAg - antibodies<br>Enzyme immunoassay kit for the qualitative and<br>quantitative determination of antibodies against HBs-<br>antigen           | 12x8 |
|        | VectoHBcAg - antibodies   |      |
| D-0566 | Enzyme immunoassay kit for the detection of total antibodies against hepatitis B core- antigen.   | 12x8 |
|        | HepaBest anti – HBc - IgG   |      |
| D-0574 | Enzyme immunoassay kit for the detection of IgG against hepatitis B core- antigen.  | 12x8 |
|        | VectoHBcAg - IgM  |      |
| D-0564 | Enzyme immunoassay kit for the detection of IgM against hepatitis B core-antigen.   | 12x8 |
|        | VectoHBe - antigen  |      |
| D-0576 | Enzyme immunoassay kit for the detection of E-antigen of the hepatitis B virus.   | 12x8 |
|        | VectoHBe - IgG  |      |
| D-0578 | Enzyme immunoassay kit for the detection of IgG against HBe-antigen of the hepatitis B virus.   | 12x8 |
|        | HEPATITIS C   |      |
|        | RecombiBest anti-HCV-IgM  |      |
| D-0760 | Enzyme immunoassay kit for the detection of IgM against hepatitis C virus.  | 12x8 |
| D-0773 | Best anti-HCV (set 1/strip)   | 24x8 |
| D-0772 | Best anti-HCV (set 2)   | 12x8 |
|        | Enzyme immunoassay kit for the detection of IgG and IgM against hepatitis C virus.  |      |
| D-0774 | Best anti-HCV – SPECTR<br>Enzyme immunoassay kit for the detection of antibodies<br>to individual proteins of hepatitis C virus (core, NS3,<br>NS4 NS5) | 6x4  |
| l      | Best anti-HCV (set 4)   |      |

| D-0776 | Enzyme immunoassay kit for the detection and confirmation of IgG and IgM against hepatitis C virus.           | 12x8 |
|--------|---|------|
| D-0777 | HCV AG/AB – IEA – BEST (set 1)  | 12x8 |
| D-0778 | HCV AG/AB – IEA – BEST (set 2)  | 24x8 |
|        | Enzyme immunoassay kit for the detection of core-<br>antigen against HCV and antibodies to hepatitis C virus. |      |
|        | HCV core-antigen-IEA – BEST   |      |
| D-0779 | Enzyme immunoassay kit for the detection of core-<br>antigen against hepatitis C virus.                       | 12x8 |
|        | HEPATITIS D   |      |
|        | Vectohep D - IgM  |      |
| D-0952 | Enzyme immunoassay kit for the detection of IgM against hepatitis D virus.                                    | 12x8 |
|        | Vectohep D - antibodies   |      |
| D-0954 | Enzyme immunoassay kit for the detection of total antibodies against hepatitis D virus.                       | 12x8 |
|        | HEPATITIS E   |      |
| D-1056 | Vectohep E - IgG<br>Enzyme immunoassay kit for the detection of IgG against<br>hepatitis E virus.             | 12x8 |
| D-1058 | Vectohep E - IgM<br>Enzyme immunoassay kit for the detection of IgM against<br>hepatitis E virus.             | 12x8 |
|        | VIRAL INTESTINAL INFECTIONS   |      |
|        | Rotavirus – antigen – EIA - BEST  |      |
| D-1652 | Enzyme immunoassay kit for the detection of human Rotavirus antigen.  | 12x8 |
| D-1654 | Adenovirus – antigen – EIA - BESTEnzyme immunoassay kit for the detection of humanAdenovirus antigen          | 12x8 |
|        | NOROVIRUS   |      |
|        | Norovirus – antigen – EIA - BEST  |      |
| D-1656 | Enzyme immunoassay kit for the detection of Norovirus antigen genogroup I and II in feces.                    | 12x8 |
|        | TICK-BORNE ENCEPHALITIS   |      |
|        | VectoTBE - IgM  |      |
| D-1152 | Enzyme immunoassay kit for the detection of IgM against tick-borne encephalitis virus.                        | 12x8 |
| D-1154 | VectoTBE - antigenEnzyme immunoassay kit for the detection of tick-borneencephalitis virus antigen.           | 12x8 |

|            | VectoTBE - IgG  |            |
|------------|---|------------|
| D-1156     | Enzyme immunoassay kit for the detection of IgG against   | 12x8       |
|            | tick-borne encephalitis virus.  |            |
|            | BORRELIOSIS   |            |
|            | LymeBest – IgG  |            |
| D-1452     | Enzyme immunoassay kit for the detection of IgG against   | 12x8       |
| D = 1 + 32 | causative agent of tick-borne borreliosis (B. afzelii, B.   | 1240       |
|            | garinii, B. burgdorferi s.s.).  | _          |
|            | LymeBest – IgM  |            |
| D-1454     | Enzyme immunoassay kit for the detection of IgM against   | 12x8       |
|            | causative agent of tick-borne borreliosis ( <i>B. afzelii, B. garinii, B. burgdorferi s.s.</i> ). |            |
| HEMOR      | RRHAGIC FEVER WITH RENAL SYND   | ROME       |
|            | (HFRS)  |            |
| D-4902     | VectoHanta – IgG  |            |
|            | Enzyme immunoassay kit for the detection of IgG against   | 12x8       |
|            | hantavirus.   |            |
| D- 4904    | VectoHanta – IgM  |            |
|            | Enzyme immunoassay kit for the detection of IgM against   | 12x8       |
|            | hantavirus.   |            |
| CRIN       | MEAN-CONGO HAEMORRHAGIC FEV   | <b>VER</b> |
|            | VectoCrimea – CHF - IgG   |            |
| D-5052     | Enzyme immunoassay kit for the detection of IgG against   | 12x8       |
|            | Crimean-Congo hemorrhagic fever virus.  |            |
|            | VectoCrimea – CHF - IgM   |            |
| D-5054     | Enzyme immunoassay kit for the detection of IgM against   | 12x8       |
|            | Crimean-Congo hemorrhagic fever virus.  |            |
|            | VectoCrimea – CHF - antigen   |            |
| D-5056     | Enzyme immunoassay kit for the detection of Crimean-  | 12x8       |
|            | Congo hemorrhagic fever virus antigen.  |            |
|            | WEST NILE VIRUS (WNV)   |            |
|            | VectoNile - IgM   |            |
| D-5150     | Enzyme immunoassay kit for the detection of IgM against   | 12x8       |
|            | West Nile virus.  |            |
|            | VectoNile - IgG   |            |
| D-5152     | Enzyme immunoassay kit for the detection of IgG against   | 12x8       |
|            | West Nile virus.  |            |
|            | VectoNile – IgG - avidity   |            |
| D-5154     | Enzyme immunoassay kit for the determination of avidity   | 6x8        |
|            | index of IgG against West Nile virus.   |            |
|            | MEASLES   |            |
|            | VectoMeasles - IgG  |            |
| D-1356     | Enzyme immunoassay kit for the detection of IgG against   | 12x8       |
|            | Measles virus.  |            |
| 1          | VectoMeasles - IgM  |            |

| D-1358 | Enzyme immunoassay kit for the detection of IgM against Measles virus.  | 12x8        |    |
|--------|---|-------------|----|
|        | MUMPS   |             |    |
| D-2602 | VectoMumps - IgG<br>Enzyme immunoassay kit for the detection of IgG against<br>Mumps virus.   | 12x8        |    |
| D-2604 | VectoMumps - IgM<br>Enzyme immunoassay kit for the detection of IgM against<br>Mumps virus.   | 12x8        |    |
|        | CYTOMEGALOVIRUS   |             |    |
| D-1552 | VectoCMV - IgM<br>Enzyme immunoassay kit for the detection of IgM against<br>Cytomegalovirus.   | 12x8        |    |
| D-1556 | CMV – IgG – EIA - BEST<br>Enzyme immunoassay kit for the quantative and<br>quantitative determination of IgG against<br>Cytomegalovirus       | 12x8        |    |
| D-1560 | <td co<="" td=""><td>20</td></td>   | <td>20</td> | 20 |
| D-1558 | VectoCMV – IgG - avidity<br>Enzyme immunoassay kit for the determination of avidity<br>index of IgG against Cytomegalovirus.                  | 6x8         |    |
| D-1566 | VectoCMV – IEA - antibodies<br>Enzyme immunoassay kit for the detection of IgG and<br>IgM against Cytomegalovirus immediate early antigen.    | 12x8        |    |
|        | TOXOPLASMA  |             |    |
| D-1752 | VectoToxo - IgG<br>Enzyme minimuoassay kit for the quantative and<br>quantitative determination of IgG against <i>Toxoplasma</i><br>gondii    | 12x8        |    |
| D-1754 | VectoToxo - IgA<br>Enzyme immunoassay kit for the detection of IgA against<br>Toxoplasma gondii.  | 12x8        |    |
|        | VectoToxo - IgM   |             |    |
| D-1756 | Enzyme immunoassay kit for the detection of IgM against <i>Toxoplasma gondii</i> .  | 12x8        |    |
| D-1760 | <b>Toxo – IgM- IEA-BEST</b><br>Enzyme immunoassay kit for the detection of IgM against<br><i>Toxoplasma gondii (</i> Antibody capture method) | 12x8        |    |
| D-1762 | VectoToxo – IgG - avidity<br>Enzyme immunoassay kit for the determination of avidity<br>index of IgG against <i>Toxoplasma gondii</i> .       | 6x8         |    |
|        | VectoToxo - antibodies  |             |    |

| D-1764 | Enzyme immunoassay kit for the detection of total antibodies against <i>Toxoplasma gondii</i> in human or  | 12x8 |
|--------|--|------|
|        | animal serum or plasma SYPHILIS  |      |
| D-1851 | RecombiBest antipallidum – IgG (set 1)   | 96x2 |
| D-1852 | RecombiBest antipallidum – IgG (set 1)<br>RecombiBest antipallidum – IgG (set 2)   | 12x8 |
| D-1632 |  | 1280 |
|        | Enzyme immunoassay kit for the detection of IgG against <i>Treponema pallidum</i> .  |      |
| D-1857 | <b>RecombiBest antipallidum - total antibodies</b> (set 1)   | 24x8 |
| D-1856 | <b>RecombiBest antipallidum - total antibodies</b> (set 2)   | 12x8 |
|        | Enzyme immunoassay kit for the detection of total antibodies against <i>Treponema pallidum</i> .   |      |
|        | RecombiBest antipallidum - IgM   |      |
| D-1858 | Enzyme immunoassay kit for the detection of IgM against <i>Treponema pallidum</i> .  | 12x8 |
|        | Treponema pallidum - blot - BEST   |      |
| D-1860 | Reagent kit for the detection of antibodies against Treponema pallidum by immunoblot.  | 24   |
|        | <b>RPHA - BEST</b> - antipallidum  |      |
| D-1820 | Reagent kit for the detection of antibodies against Treponema pallidum by passive hemagglutination test.   | 100  |
|        |  |      |
|        | CHLAMYDIOSIS   |      |
|        | Chlamydophila pneumoniae – IgG – EIA - BEST  |      |
| D-1944 | Enzyme immunoassay kit for the detection of IgG against <i>Chlamydophila pneumoniae</i> .  | 12x8 |
|        | Chlamydophila pneumoniae – IgM – EIA - BEST  |      |
| D-1946 | Enzyme immunoassay kit for the detection of IgM against <i>Chlamydophila pneumoniae</i> .  | 12x8 |
|        | Chlamydophila pneumoniae – IgA – EIA - BEST  |      |
| D-1948 | Enzyme immunoassay kit for the detection of IgA against <i>Chlamydophila pneumoniae</i> .  | 12x8 |
| D-1964 | ChlamyBest C.trachomatis - IgG           Enzyme immunoassay kit for the detection of IgG against           Chlamydia trachomatis .                   | 12x8 |
|        | ChlamyBest C.trachomatis - IgM   |      |
| D-1966 | Enzyme immunoassay kit for the detection of IgM against <i>Chlamydia trachomatis</i> .   | 12x8 |
| D-1968 | ChlamyBest C.trachomatis - IgA           Enzyme immunoassay kit for the detection of IgA against           Chlamydia trachomatis .                   | 12x8 |
| D-1972 | ChlamyBest cHSP60 - IgGEnzyme immunoassay kit for the detection of IgG againstChlamydia trachomatis heat shock protein (cHSP60).                     | 12x8 |
| D-1982 | ChlamyBest MOMP + pgp3 - IgG<br>Enzyme immunoassay kit for the detection of IgG against<br><i>Chlamydia trachomatis</i> major outer membrane protein | 12x8 |

| TRICHOMONIASIS |   |      |
|----------------|---|------|
|                | Trichomonas vaginalis - IgG – EIA - BEST  |      |
| D-2052         | Enzyme immunoassay kit for the detection of IgG against <i>Trichomonas vaginalis</i> .              | 12x8 |
|                | Trichomonas vaginalis – IgA – EIA - BEST  |      |
| D-2058         | Enzyme immunoassay kit for the detection of IgA against   | 12x8 |
|                | Trichomonas vaginalis.  |      |
|                | HERPES VIRUSES  |      |
|                | VectoHSV - IgG  |      |
| D-2152         | Enzyme immunoassay kit for the detection of IgG against   | 12x8 |
|                | herpes simplex virus.   |      |
|                | VectoHSV - 1,2 – IgG - avidity  |      |
| D-2156         | Enzyme immunoassay kit for the determination of avidity   | 6x8  |
|                | index of IgG against HSV-1, 2.  |      |
|                | VectoHSV - IgM  |      |
| D-2154         | Enzyme immunoassay kit for the detection of IgM against   | 12x8 |
|                | HSV-1, 2.   |      |
|                | VectoHSV-2 - IgG  |      |
| D-2180         | Enzyme immunoassay kit for the detection of IgG against   | 12x8 |
|                | HSV-2.  |      |
| D 0100         | VectoHSV-2 – IgG - avidity  | 6.0  |
| D-2182         | Enzyme immunoassay kit for the determination of avidity   | 6x8  |
|                | index of IgG against HSV-2.   |      |
| D 2101         | VectoHSV-2 - IgM  | 12x8 |
| D-2181         | Enzyme immunoassay kit for the detection of IgM against HSV-2.                                      | 1288 |
|                | VectoVZV - IgG  |      |
| D-2192         | Enzyme immunoassay kit for the detection of IgG against   | 12x8 |
|                | Varicella-Zoster virus.   |      |
|                | VectoVZV - IgM  |      |
| D-2188         | Enzyme immunoassay kit for the detection of IgM against   | 12x8 |
|                | Varicella-Zoster virus.   |      |
|                | VectoVZV – gE - IgG   |      |
| D-2186         | Enzyme immunoassay kit for the detection of IgG against   | 12x8 |
|                | glycoprotein E of Varicella-Zoster virus.   |      |
|                | VectoEBV – VCA - IgG  |      |
| D-2184         | Enzyme immunoassay kit for the detection of IgG against   | 12x8 |
|                | viral capsid antigen of Epstein-Barr virus.   |      |
| D 0100         | VectoEBV–VCA–IgG –avidity   | 6.0  |
| D-2183         | Enzyme immunoassay kit for the determination of IgG   | 6x8  |
|                | avidity index against VCA of EBV.   |      |
| D 2176         | VectoEBV – VCA - IgM  | 12-0 |
| D-2176         | Enzyme immunoassay kit for the detection of IgM against viral capsid antigen of Epstein-Barr virus. | 12x8 |
|                | vitai capsia antigen of Epstein-Dall vitus.   |      |

|         | VectoEBV – NA - IgG                                     |       |
|---------|---|-------|
| D-2170  | Enzyme immunoassay kit for the detection of IgG against | 12x8  |
|         | nuclear antigen (NA) of Epstein-Barr virus.             |       |
|         | VectoEBV – EA - IgG                                     |       |
| D-2172  | Enzyme immunoassay kit for the detection of IgG against | 12x8  |
|         | early antigen (EA) of Epstein-Barr virus.               |       |
|         | VectoHHV-6 - IgG  |       |
| D-2166  | Enzyme immunoassay kit for the detection of IgG against | 12x8  |
| D 2100  | HHV-6.  | 12/10 |
|         | VectoHHV-8 – IgG  |       |
| D-2160  | Enzyme immunoassay kit for the detection of IgG against | 12x8  |
|         | HHV-8.  |       |
|         | UREAPLASMOSIS   |       |
|         | Ureaplasma urealyticum – IgG – EIA - BEST               |       |
| D-2254  | Enzyme immunoassay kit for the detection of IgG against | 12x8  |
|         | Ureaplasma urealyticum antigens.                        |       |
|         | Ureaplasma urealyticum – IgA – EIA - BEST               |       |
| D-2258  | Enzyme immunoassay kit for the detection of IgA against | 12x8  |
|         | Ureaplasma urealyticum antigens.                        |       |
|         | MYCOPLASMOSIS   |       |
|         | Mycoplasma hominis – IgG – EIA - BEST                   |       |
| D-4352  | Enzyme immunoassay kit for the detection of IgG against | 12x8  |
|         | Mycoplasma hominis .                                    |       |
|         | Mycoplasma hominis – IgA – EIA - BEST                   |       |
| D-4358  | Enzyme immunoassay kit for the detection of IgA against | 12x8  |
|         | Mycoplasma hominis .                                    |       |
|         | Mycoplasma pneumonia – IgG – EIA - BEST                 |       |
| D-4362  | Enzyme immunoassay kit for the detection of IgG against | 12x8  |
|         | Mycoplasma pneumoniae .                                 |       |
|         | Mycoplasma pneumonia – IgM – EIA - BEST                 |       |
| D-4366  | Enzyme immunoassay kit for the detection of IgM against | 12x8  |
|         | Mycoplasma pneumoniae.                                  |       |
| D 10 (0 | Mycoplasma pneumonia – IgA – EIA - BEST                 |       |
| D-4368  | Enzyme immunoassay kit for the detection of IgA against | 12x8  |
|         | Mycoplasma pneumoniae .                                 |       |
|         | TUBERCULOSIS  |       |
|         | AB – Tub - BEST   |       |
| D-2352  | Enzyme immunoassay kit for the detection of total       | 12x8  |
|         | antibodies against <i>Mycobacterium tuberculosis</i> .  |       |
|         | BRUCELLOSIS   |       |
|         | Brucellosis – IgG – EIA - BEST                          |       |
| D-3652  | Enzyme immunoassay kit for the detection of IgG against | 12x8  |
|         | Brucellosis.  |       |
|         | Brucellosis – IgA – EIA - BEST                          | 1.    |
| D-3654  | Enzyme immunoassay kit for the detection of IgA against | 12x8  |
|         | Brucellosis.  |       |

|        | Brucellosis – IgM – EIA - BEST                          |      |
|--------|---|------|
| D-3656 | Enzyme immunoassay kit for the detection of IgM against | 12x8 |
|        | Brucellosis.  |      |
| D-3658 | Brucellosis – antibodies – EIA - BEST                   |      |
|        | Enzyme immunoassay kit for the detection of total       | 12x8 |
|        | antibodies against Brucellosis.                         |      |
|        | YERSINIOSIS   |      |
|        | Yersinia – IgG – EIA –BEST                              |      |
| D-3202 | Enzyme immunoassay kit for the detection of IgG against | 12x8 |
|        | Yersinia.   |      |
|        | Yersinia – IgA – EIA –BEST                              |      |
| D-3204 | Enzyme immunoassay kit for the detection of IgA against | 12x8 |
|        | Yersinia.   |      |
|        | Yersinia – IgM – EIA –BEST                              |      |
| D-3206 | Enzyme immunoassay kit for the detection of IgG against | 12x8 |
|        | Yersinia.   |      |
|        | RUBELLA   |      |
|        | VectoRubella - IgG                                      |      |
| D-2552 | Enzyme immunoassay kit for the detection of IgG against | 12x8 |
|        | Rubella virus.  |      |
|        | VectoRubella - IgM                                      |      |
| D-2554 | Enzyme immunoassay kit for the detection of IgM against | 12x8 |
|        | Rubella virus.  |      |
|        | VectoRubella – IgG - avidity                            |      |
| D-2556 | Enzyme immunoassay kit for the determination of avidity | 6x8  |
|        | index of IgG against Rubella virus.                     |      |
|        | Rubella – IgM-EIA-BEST                                  |      |
| D-2560 | Enzyme immunoassay kit for the detection of IgM to      | 12x8 |
|        | Rubella virus by «capture» method.                      |      |
|        | PARASITIC DISEASES                                      |      |
|        | Toxocara – IgG – EIA - BEST                             |      |
| D-2752 | Enzyme immunoassay kit for the detection of IgG against | 12x8 |
|        | Toxocara canis antigens.                                |      |
|        | Opisthorchis – IgG – EIA - BEST                         |      |
| D-2952 | Enzyme immunoassay kit for the detection of IgG against | 12x8 |
|        | Opisthorchis felineus antigens.                         |      |
|        | <b>Opisthorchis – IgM – EIA - BEST</b>                  |      |
| D-2954 | Enzyme immunoassay kit for the detection of IgM against | 12x8 |
|        | Opisthorchis felineus antigens.                         |      |
|        | <b>Opisthorchis – CIC – EIA - BEST</b>                  |      |
| D-2956 | Enzyme immunoassay kit for the detection of specific    | 12x8 |
|        | circulatory immune complexes (CIC), containing          |      |
|        | Clonorchis – IgG – EIA - BEST                           |      |
| D-2958 | Enzyme immunoassay kit for the detection of IgG against | 12x8 |
|        | Clonorchis sinensis antigens.                           |      |
|        | Trichinella – IgG – EIA - BEST                          |      |
| D-3152 | Enzyme immunoassay kit for the detection of IgG against | 12x8 |
|        | Trichinella spiralis antigens.                          |      |

|                  | Trichinella – IgM – EIA - BEST  |             |
|------------------|---|-------------|
| D-3154           | Enzyme immunoassay kit for the detection of IgM against   | 12x8        |
|                  | Trichinella spiralis antigens.  |             |
|                  | Helminths – IgG – EIA – BEST  |             |
| D-3354           | Enzyme immunoassay kit for the detection of IgG against   | 3x64        |
|                  | Opisthorchis felineus, Trichinella, Toxocara and  |             |
|                  | Echinococcus – IgG – EIA - BEST   |             |
| D-3356           | Enzyme immunoassay kit for the detection of IgG against   | 12x8        |
|                  | Echinococcus granulosus antigens.   |             |
|                  | Ascaris – IgG – EIA - BEST  |             |
| D-3452           | Enzyme immunoassay kit for the detection of IgG against   | 12x8        |
|                  | Ascaris lumbricoides antigens.  |             |
|                  | Anisakis-IgG-EIA-BEST   |             |
| D-3454           | Enzyme immunoassay kit for the detection of IgG against   | 12x8        |
|                  | Anisakis nematodes antigens.  |             |
|                  | Cysticercus-IgG-EIA-BEST  |             |
| D-3456           | Enzyme immunoassay kit for the detection of IgG against   | 12x8        |
|                  | Taenia solium antigens.   |             |
|                  | Lamblia – antibodies – EIA - BEST   |             |
| D-3552           | Enzyme immunoassay kit for the detection of IgA, IgM  | 12x8        |
|                  | and IgG against Giardia lamblia antigens.   |             |
|                  | Lamblia – IgM – EIA - BEST  |             |
| D-3554           | Enzyme immunoassay kit for the detection of IgM against   | 12x8        |
|                  | Giardia lamblia antigens.   |             |
|                  | Lamblia – antigen – EIA - BEST  |             |
| D-3556           |   | 12x8        |
| 2 0000           | lamblia antigens in faeces.   |             |
| (                | GASTROINTESTINAL DISEASES   |             |
|                  | HelicoBest - antibodies   |             |
| D-3752           | Enzyme immunoassay kit for the detection of total   | 12x8        |
|                  | antibodies against CagA antigen of <i>Helicobacter pylori</i> .   | 12/10       |
|                  | IgG – Gliadin – EIA - BEST  |             |
|                  | Enzyme immunoassay kit for the quantitative   |             |
| D-3754           | determination of IgG against gliadin.   | 12x8        |
|                  | Measurement range: 0-100 IU/ml.   | -           |
|                  | Sensitivity: 2.5 IU/ml.   |             |
|                  | IgA – Gliadin – EIA - BEST  |             |
|                  | Enzyme immunoassay kit for the quantitative   |             |
| D-3756           | determination of IoA accinetation   |             |
|                  | determination of IgA against gliadin.   | 12x8        |
|                  | determination of IgA against gliadin.<br>Measurement range: 0-100 IU/ml.  | 12x8        |
|                  | Measurement range: 0-100 IU/ml.   | 12x8        |
|                  | Measurement range: 0-100 IU/ml.<br>Sensitivity: 2.5 IU/ml.  | 12x8        |
|                  | Measurement range: 0-100 IU/ml.   | 12x8        |
| D-3758           | Measurement range: 0-100 IU/ml.<br>Sensitivity: 2.5 IU/ml.<br>IgA – transglutaminase – EIA - BEST<br>Enzyme immunoassay kit for the quantitative  | 12x8<br>6x8 |
| D-3758           | Measurement range: 0-100 IU/ml.<br>Sensitivity: 2.5 IU/ml.<br>IgA – transglutaminase – EIA - BEST   |             |
| D-3758           | Measurement range: 0-100 IU/ml.Sensitivity: 2.5 IU/ml.IgA – transglutaminase – EIA - BESTEnzyme immunoassay kit for the quantitativedetermination of IgA auto-antibodies against tissue   |             |
| D-3758           | Measurement range: 0-100 IU/ml.<br>Sensitivity: 2.5 IU/ml.<br><b>IgA – transglutaminase – EIA - BEST</b><br>Enzyme immunoassay kit for the quantitative<br>determination of IgA auto-antibodies against tissue<br>Measurement range: 2.5-100 IU/ml. |             |
| D-3758<br>D-3760 | Measurement range: 0-100 IU/ml.Sensitivity: 2.5 IU/ml.IgA – transglutaminase – EIA - BESTEnzyme immunoassay kit for the quantitativedetermination of IgA auto-antibodies against tissueMeasurement range: 2.5-100 IU/ml.Sensitivity: 2.5 IU/ml.     | 6x8         |

|        | Pepsinogen I – EIA - BEST                                |      |
|--------|--|------|
|        | Enzyme immunoassay kit for the quantitative              |      |
| D-3762 | determination of Pepsinogen I.                           | 12x8 |
|        | Measurement range: 0-200 mkg/l.                          |      |
|        | Sensitivity: 1 mkg/l.                                    |      |
|        | Pepsinogen II – EIA - BEST                               |      |
|        | Enzyme immunoassay kit for the quantitative              |      |
| D-3764 | determination of Pepsinogen II.                          | 12x8 |
|        | Measurement range: 0-50 mkg/l.                           |      |
|        | Sensitivity: 1 mkg/l.                                    |      |
|        | HORMONES   |      |
|        | TSH – EIA - BEST   |      |
|        | Enzyme immunoassay kit for the quantitative              |      |
| X-3952 | determination of thyroid stimulation hormone (TSH).      | 12x8 |
|        | Measurement range: 0-16 mIU/l.                           |      |
|        | Sensitivity: 0.1 mIU/l.                                  |      |
|        | T <sub>3</sub> total – EIA - BEST                        |      |
|        | Enzyme immunoassay kit for the quantitative              |      |
| X-3954 | determination of total triiodothyronine.                 | 12x8 |
|        | Measurement range: 0-9 nmol/l.                           |      |
|        | Sensitivity: 0.2 nmol/l.                                 |      |
|        | T <sub>3</sub> free – EIA - BEST                         |      |
|        | Enzyme immunoassay kit for the quantitative              |      |
| X-3970 | determination of free triiodothyronine.                  | 12x8 |
|        | Measurement range: 0-20 pmol/l.                          |      |
|        | Sensitivity: 0.5 pmol/l.                                 |      |
|        | T <sub>4</sub> total – EIA - BEST                        |      |
|        | Enzyme immunoassay kit for the quantitative              |      |
| X-3956 | determination of total thyroxin.                         | 12x8 |
|        | Measurement range: 0-400 nmol/l.                         |      |
|        | Sensitivity: 10 nmol/l.                                  |      |
|        | T <sub>4</sub> free – EIA - BEST                         |      |
|        | Enzyme immunoassay kit for the quantitative              |      |
| X-3962 | determination of free thyroxin.                          | 12x8 |
|        | Measurement range: 0-80 pmol/l.                          |      |
|        | Sensitivity: 1 pmol/l.                                   |      |
|        | TG – EIA - BEST  |      |
| X-3966 | Enzyme immunoassay kit for the quantitative              | 12x8 |
|        | determination of thyroglobulin.                          |      |
|        | Anti-TG – EIA - BEST                                     |      |
|        | Enzyme immunoassay kit for the quantitative              |      |
| X-3958 | determination of total antibodies against thyroglobulin. | 12x8 |
|        | Measurement range: 0-2000 IU/ml.                         |      |
|        | Sensitivity: 10 IU/ml.                                   |      |
|        | Anti – TPO – EIA - BEST                                  |      |
|        | Enzyme immunoassay kit for the quantitative              |      |
| X-3968 | determination of total antibodies against thyroid        | 12x8 |
|        | Measurement range: 0-1000 IU/ml.                         |      |
|        | Sensitivity: 5 IU/ml.                                    |      |

|        | Prolactin – EIA – BEST                                    |      |
|--------|---|------|
|        | Enzyme immunoassay kit for the quantitative               |      |
| X-3960 | determination of prolactin.                               | 12x8 |
|        | Measurement range: 0-3000 mIU/l.                          |      |
|        | Sensitivity: 15 mIU/l.                                    |      |
|        | Cortisol – EIA - BEST                                     |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| X-3964 | determination of cortisol.                                | 12x8 |
|        | Measurement range: 0-1200 nmol/l                          |      |
|        | Sensitivity: 5 nmol/l.                                    |      |
|        | Testosterone – EIA - BEST                                 |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| X-3972 | determination of testosterone.                            | 12x8 |
|        | Measurement range: 0-60 nmol/l.                           |      |
|        | Sensitivity: 0.2 nmol/l.                                  |      |
|        | FSH – EIA - BEST  |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| X-3974 | determination of follicle-stimulating hormone.            | 12x8 |
|        | Measurement range: 0-100 mIU/ml.                          |      |
|        | Sensitivity: 1 mIU/ml.                                    |      |
|        | LH – EIA - BEST   |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| X-3976 | determination of luteinizing hormone.                     | 12x8 |
|        | Measurement range: 0-100 mIU/ml.                          |      |
|        | Sensitivity: 0.5 mIU/ml.                                  |      |
|        | Progesterone – EIA - BEST                                 |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| X-3978 | determination of progesterone.                            | 12x8 |
|        | Measurement range: 0-100 nmol/l.                          |      |
|        | Sensitivity: 0.4 nmol/l.                                  |      |
|        | PREGNANCY   |      |
|        | HCG – EIA - BEST  |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| D-4154 | determination of human chorionic gonadotropin.            | 12x8 |
|        | Measurement range: 0-500 MIU/l.                           |      |
|        | Sensitivity: 2 MIU/l.                                     |      |
|        | Free β-HCG – EIA - BEST                                   |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| D-4158 | determination of free $\beta$ -subunit of human chorionic | 12x8 |
|        | Measurement range: 0-200 ng/ml.                           |      |
|        | Sensitivity: 1 ng/ml.                                     |      |
|        | PAPP-A – EIA - BEST                                       |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| D-4160 | determination of pregnancy-associated plasma protein A    | 12x8 |
|        | Measurement range: 0-10000 mIU/l.                         |      |
|        | Sensitivity: 20 mIU/l.                                    |      |
|        | Prolactin – EIA - BEST                                    |      |

|        | Enzyme immunoassay kit for the quantitative  |      |
|--------|--|------|
| X-3960 | determination of prolactin.  | 12x8 |
|        | Measurement range: 0-3000 mIU/l.   |      |
|        | Sensitivity: 15 mIU/l.   |      |
|        | AFP – EIA - BEST   |      |
|        | Enzyme immunoassay kit for the quantitative  |      |
| T-8456 | determination of   | 12x8 |
| 1 0450 | α-Fetoprotein.   |      |
|        | Measurement range: 0-400 IU/ml.  |      |
|        | Sensitivity: 2.5 IU/ml.  |      |
|        | TBG – EIA - BEST   |      |
|        | Enzyme immunoassay kit for the quantitative  |      |
| T-8468 | determination of trophoblastic $\beta$ -1-glycoprotein.  | 12x8 |
|        | Measurement range: 0-200 ng/ml.  |      |
|        | Sensitivity: 1.5 ng/ml.  |      |
|        | Ferritin – EIA - BEST  |      |
|        | Enzyme immunoassay kit for the quantitative  |      |
| T 9557 | determination of trophoblastic $\beta$ -1-glycoprotein.  | 120  |
| T-8552 | ferritin.  | 12x8 |
|        | Measurement range: 0-500 ng/ml.  |      |
|        | Sensitivity: 2.5 ng/ml.  |      |
|        | CANDIDIASIS  |      |
|        | Candida – IgG – EIA - BEST   | 12x8 |
| D-4652 | Enzyme immunoassay kit for the detection of IgG against  |      |
|        | Candida fungi.   |      |
|        | Candida – IgM – EIA - BEST   |      |
| D-4654 | Enzyme immunoassay kit for the detection of IgM against  | 12x8 |
|        | Candida fungi.   |      |
|        | Candida – IgA – EIA - BEST   | 10 0 |
| D-4656 | Enzyme immunoassay kit for the detection of IgA against  | 12x8 |
|        | Candida fungi.   |      |
|        | ASPERGILLOSIS  |      |
|        | Aspergill – IgG – EIA - BEST   |      |
| D-4752 | Enzyme immunoassay kit for the detection of IgG against  | 12x8 |
|        | Aspergillus fungi.   |      |
|        | ANAEMIA  |      |
|        | Ferritin – EIA - BEST  |      |
|        | Enzyme immunoassay kit for the quantitative  |      |
| Т-8552 | determination of ferritin.   | 12x8 |
|        | Measurement range: 0-500 ng/ml.  |      |
|        |  |      |
|        |  |      |
| A-8776 | 5 5 1  |      |
|        |  | 12x8 |
|        |  |      |
|        | Sensitivity: 0,5 mIU/ml.   |      |
| A-8776 | Measurement range: 0-500 ng/ml.         Sensitivity: 2,5 ng/ml.         Erythropoietin – EIA - BEST         Enzyme immunoassay kit for the quantitative         determination of erythropoietin.         Measurement range: 0-200 mIU/ml.         Sensitivity: 0,5 mIU/ml.         TUMOR MARKERS | 12x8 |

|        | HCG – EIA - BEST   |        |
|--------|--|--------|
|        | Enzyme immunoassay kit for the quantitative                              |        |
| D-4154 | determination of human chorionic gonadotropin.                           | 12x8   |
|        | Measurement range: 0-500 IU/l.   | 12.110 |
|        | Sensitivity: 2 IU/l.   |        |
|        | 5  |        |
|        | Free β-HCG – EIA - BEST<br>Enzyme minimunoassay και τοι της quantitative |        |
| D-4158 | determination of free $\beta$ -subunit of human chorionic                | 12x8   |
|        | Measurement range: 0-200 ng/ml.  | 1270   |
|        | Sensitivity: 1 ng/ml.  |        |
|        | CEA – EIA - BEST   |        |
|        | Enzyme immunoassay kit for the quantitative                              |        |
| T-8454 | determination of carcinoembryonal antigen.                               | 12x8   |
|        | Measurement range: 0-80 ng/ml.   | 12/10  |
|        | Sensitivity: 0.3 ng/ml.  |        |
|        | AFP - EIA - BEST   |        |
|        | Enzyme immunoassay kit for the quantitative                              |        |
| T-8456 | determination of $\alpha$ -fetoprotein.                                  | 12x8   |
|        | Measurement range: 0-400 IU/ml.  |        |
|        | Sensitivity: 2.5 IU/ml.  |        |
|        | PSA total – EIA - BEST   |        |
|        | Enzyme immunoassay kit for the quantitative                              |        |
| T-8453 | determination of total prostate specific antigen (PSA).                  | 12x8   |
|        | Measurement range: 0-5 ng/ml.  |        |
|        | Sensitivity: 0.01 ng/ml.   |        |
|        | PSA free – EIA - BEST  | 12x8   |
|        | Enzyme immunoassay kit for the quantitative                              |        |
| T-8460 | determination of free prostate specific antigen (PSA).                   |        |
|        | Measurement range: 0-5 ng/ml.  |        |
|        | Sensitivity: 0.1 ng/ml.  |        |
|        | CA – 125 – EIA - BEST  |        |
|        | Enzyme immunoassay kit for the quantitative                              |        |
| T-8466 | determination of CA-125.   | 12x8   |
|        | Measurement range: 0-400 IU/ml.  |        |
|        | Sensitivity: 1.5 IU/ml.  |        |
|        | TPBG – EIA - BEST  |        |
|        | Enzyme immunoassay kit for the quantitative                              | 12x8   |
| T-8468 | determination of trophoblastic $\beta$ -1-glycoprotein.                  |        |
|        | Measurement range: 0-200 ng/ml.  |        |
|        | Sensitivity: 1.5 ng/ml.  |        |
|        | CA 19-9 – EIA - BEST   | 12x8   |
| T-8470 | Enzyme immunoassay kit for the quantitative                              |        |
| 1 01/0 | Measurement range: 0-300 IU/ml.  |        |
|        | Sensitivity: 2 IU/ml.  |        |
|        | CA 15-3 – EIA - BEST   | 12x8   |
| T-8472 | Enzyme immunoassay kit for the quantitative                              |        |
| 1 0174 | Measurement range: 0-250 IU/ml.  |        |
|        | Sensitivity: 1,5 IU/ml.  |        |
|        | Ferritin-EIA-BEST  |        |

|        | Enzyme immunoassay kit for the quantitative               |      |
|--------|---|------|
| T-8552 | determination of human ferritin.                          | 12x8 |
|        | Measurement range: 0-500 ng/ml                            |      |
|        | Sensitivity: 2.5 ng/ml.                                   |      |
|        | Pepsinogen I – EIA - BEST                                 |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| D-3762 | determination of pepsinogen I.                            | 12x8 |
|        | Measurement range: 0-200 mkg/l.                           |      |
|        | Sensitivity: 1 mkg/l.                                     |      |
|        | Pepsinogen II – EIA - BEST                                |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| D-3764 | determination of pepsinogen II.                           | 12x8 |
|        | Measurement range: 0-50 mkg/l.                            |      |
|        | Sensitivity: 1 mkg/l.                                     |      |
|        | ACUTE PHASE PROTEINS                                      |      |
|        | CRP – EIA – BEST  |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| A-9002 | determination of C-reactive protein (CRP)                 | 12x8 |
|        | Measurement range: 0-10 IU/l                              |      |
|        | Sensitivity: 0.05 IU/l                                    |      |
|        | Procalcitonin – EIA –BEST                                 |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| A-9004 | determination of Procalcitonin.                           | 12x8 |
|        | Measurement range: 0-12,8 ng/ml.                          |      |
|        | Sensitivity: 0,04 ng/ml.                                  |      |
|        | CARDIAC MARKERS   |      |
|        | CRP – EIA – BEST  |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| A-9002 | determination of C-reactive protein (CRP).                | 12x8 |
|        | Measurement range: 0-10 IU/l.                             |      |
|        | Sensitivity: 0.05 IU/l.                                   |      |
|        | NTproBNP – EIA - BEST                                     |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| A-9102 | determination of N-terminal fragment of brain natriuretic | 12x8 |
|        | Measurement range: 0-2000 pg/ml.                          |      |
|        | Sensitivity: 20 pg/ml.                                    |      |
|        | FABP – EIA - BEST   |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| A-9104 | determination of fatty-acid-binding protein.              | 12x8 |
|        | Measurement range: 0-15 ng/ml.                            |      |
|        | Sensitivity: 0.05 ng/ml.                                  |      |
|        | Troponin I – EIA - BEST                                   |      |
|        | Enzyme immunoassay kit for the quantitative               |      |
| A-9106 | determination of troponin I.                              | 12x8 |
|        | Measurement range: 0-6 ng/ml.                             | 1240 |
|        | Sensitivity: 0.02 ng/ml.                                  |      |
|        |   |      |

| A-9108  | Enzyme immunoassay kit for the quantitative determination of myoglobin.               | 12x8    |
|---------|---|---------|
|         | Measurement range: 0-1000 ng/ml.  |         |
|         | Sensitivity: 4 ng/ml.   |         |
|         | TROMBOSIS   |         |
|         | D-dimer - EIA – BEST  |         |
| D-9120  | Enzyme immunoassay kit for the quantitative determination of D-dimer in human plasma. | 12x8    |
|         | Measurement range: 0-1500 ng/ml   |         |
| QVC     | Sensitivity: 10 ng/ml   | 20      |
| 513     | STEMIC AND AUTOIMMUNE DISEASE   | 20      |
|         | <b>IgG – Gliadin – EIA - BEST</b><br>Enzyme immunoassay kit for the quantitative      |         |
| D-3754  | determination of IgG against gliadin.   | 12x8    |
| D-3734  | Measurement range: 0-100 IU/ml.   | 1280    |
|         | Sensitivity: 2.5 IU/ml.   |         |
|         | IgA – Gliadin – EIA - BEST  |         |
|         | Enzyme immunoassay kit for the quantitative   |         |
| D-3756  | determination of IgA against gliadin.   | 12x8    |
| D-3730  | Measurement range: 0-100 IU/ml.   | 1240    |
|         | Sensitivity: 2.5 IU/ml.   |         |
|         | IgA – transglutaminase – EIA - BEST   |         |
|         | Enzyme immunoassay kit for the quantitative   |         |
| D-3758  | determination of IgA auto-antibodies against tissue                                   | 6x16    |
| D 5750  | Measurement range: 2.5-100 U/ml.  |         |
|         | Sensitivity: 2.5 U/ml.  |         |
|         | Anti – TG – EIA - BEST  |         |
|         | Enzyme immunoassay kit for the quantitative   |         |
| X-3958  | determination of total antibodies against thyreoglobulin.                             | 12x8    |
|         | Measurement range: 0-2000 IU/ml.  |         |
|         | Sensitivity: 10 IU/ml.  |         |
|         | Anti – TPO – EIA – BEST   |         |
|         | Enzyme immunoassay kit for the quantitative   |         |
| X-3968  | determination of total antibodies against thyroid                                     | 12x8    |
|         | Measurement range: 0-1000 IU/ml.  |         |
|         | Sensitivity: 5 IU/ml.   |         |
|         | VectoRheumatoid factor M-class  |         |
| A-8652  | Enzyme immunoassay kit for the quantitative   | 12x8    |
|         | determination of M-class Rheumatoid factor.   | - = 110 |
|         | VectoRheumatoid factor total  | ·       |
| A-8654  | Enzyme immunoassay kit for the quantitative   | 12x8    |
|         | determination of total Rheumatoid factor.   | 12/10   |
|         | Vecto-dsDNA–IgG   |         |
| A-8656  | Enzyme immunoassay kit for the quantitative determination                             | 12x8    |
| 41-0030 | of IgG autoimmune antibodies against double-stranded DNA.                             | 12110   |
|         | Vecto - ssDNA – IgG   |         |
| A-8658  | Enzyme immunoassay kit for the quantitative determination                             | 12x8    |
|         | of IgG autoimmune antibodies against single-stranded DNA.                             |         |

|         | IgE total – EIA – BEST   |       |
|---------|--|-------|
|         | Enzyme immunoassay kit for the quantitative  |       |
| A-8660  | determination of total IgE.  | 12x8  |
| 11 0000 | Measurement range: 0-800 IU/ml.  | 12/10 |
|         | Sensitivity: 2.5 IU/ml.  |       |
| CONTD   |  | ATIC  |
| CONTR   | OL OF THE HUMORAL IMMUNE ST<br>IgE total – EIA - BEST                                  | AIUS  |
|         | Enzyme immunoassay kit for the quantitative  |       |
| A-8660  | determination of total IgE.  | 12x8  |
| A-0000  | Measurement range: 0-800 IU/ml.  | 1280  |
|         | Sensitivity: 2.5 IU/ml.  |       |
|         | IgG total – EIA - BEST   |       |
|         | Enzyme immunoassay kit for the quantitative  |       |
| A-8662  | Measurement range: 0-24 mg/ml.   | 12x8  |
|         | Sensitivity: 0.2 mg/ml.  |       |
|         | IgM total – EIA - BEST   |       |
|         | <u> </u>   |       |
| A-8664  | Enzyme immunoassay kit for the quantitative  | 12x8  |
|         | Measurement range: 0-3.2 mg/ml.  |       |
|         | Sensitivity: 0.032 mg/ml.  |       |
|         | IgA total – EIA - BEST   |       |
| A-8666  | Enzyme immunoassay kit for the quantitative  | 12x8  |
|         | Measurement range: 0-4.2 mg/ml.  |       |
|         | Sensitivity: 0.021 mg/ml.  |       |
| A 0/74  | Immunoscreen G, M, A – EIA - BEST  | 22.2  |
| A-8674  | Enzyme immunoassay kit for the determination of total                                  | 32x3  |
|         | IgA, IgM, IgG concentration.   |       |
|         | <b>IgA secretory – EIA - BEST</b><br>Enzyme immunoassay kit for the quantitative       |       |
| A-8668  | determination of secretory IgA (sIgA).   | 12x8  |
| A-8008  |  | 1288  |
|         | Measurement range: 0-20 mg/l.  |       |
|         | Sensitivity: 0.35 mg/l.  |       |
|         | CYTOKINES  |       |
|         | γ-Interferon – EIA - BEST  |       |
| A 9753  | Enzyme immunoassay kit for the quantitative  | 120   |
| A-8752  | determination of $\gamma$ -interferon in human biological fluids                       | 12x8  |
|         | Measurement range: 0-1000 pg/ml.   |       |
|         | Sensitivity: 5 pg/ml.  |       |
|         | Interleukin-4 – EIA - BEST   |       |
| A 9751  | Enzyme immunoassay kit for the quantitative  | 12x8  |
| A-8754  | determination of interleukin-4 in human biological fluids                              |       |
|         | Measurement range: 0-100 pg/ml.  |       |
|         | Sensitivity: 0.4 pg/ml.  |       |
|         | a -TNF - EIA-BEST  |       |
| A 0757  | Enzyme immunoassay kit for the quantitative  | 100   |
| A-8756  | determination of tumor necrosis factor - α in human<br>Measurement range: 0-250 pg/ml. | 12x8  |
|         | $\mu_{\nu}$  |       |
|         | Sensitivity: 2 pg/ml.  |       |

|         | Enzyme immunoassay kit for the quantitative                               |      |
|---------|---|------|
| A-8758  | determination of $\alpha$ -interferon in human biological fluids          | 12x8 |
|         | Measurement range: 0-500 pg/ml.   |      |
|         | Sensitivity: 5 pg/ml.   |      |
|         | α-Interferon - autoimmune antibodies – EIA - BEST                         |      |
|         | Enzyme immunoassay kit for the quantitative                               |      |
| A-8760  | determination of autoimmune antibodies against $\alpha$ -                 | 12x8 |
| 11 0700 | Measurement range: 0-100 ng/ml.   | 12/0 |
|         | Sensitivity: 0.4 ng/ml.   |      |
|         | Interleukin-8 – EIA - BEST  |      |
|         | Enzyme immunoassay kit for the quantitative                               |      |
| A-8762  | determination of interleukin-8 in human biological fluids                 | 12x8 |
| 11-0702 | Measurement range: 0-250 pg/ml.   | 1240 |
|         | Sensitivity: 2 pg/ml.   |      |
|         | Interleukin-1 β – EIA - BEST  |      |
|         | Enzyme immunoassay kit for the quantitative                               |      |
| A-8766  | determination of interleukin-1 beta in human biological                   | 12x8 |
| A-0/00  | Measurement range: 0-250 pg/ml.   | 1280 |
|         | Sensitivity: 1 pg/ml.   |      |
|         |   |      |
|         | Interleukin-6 – EIA - BEST<br>Enzyme minunoassay kit for the quantitative |      |
|         | determination of interleukin-6 in human biological fluids                 | 10 0 |
| A-8768  | and culture medium  | 12x8 |
|         | Measurement range: 0-300 pg/ml.   |      |
|         | Sensitivity: 0.5 pg/ml.   |      |
|         | Interleukin-18 – EIA - BEST   |      |
|         | Enzyme immunoassay kit for the quantitative                               |      |
| A-8770  | determination of interleukin-18 in human biological                       | 12x8 |
|         | Measurement range: 0-1000 pg/ml.  |      |
|         | Sensitivity: 5 pg/ml.   |      |
|         | Interleukin-2 – EIA - BEST  |      |
|         | Enzyme immunoassay kit for the quantitative                               |      |
| A-8772  | determination of interleukin-2 in human biological fluids                 | 12x8 |
|         | Measurement range: 0-500 pg/ml.   |      |
|         | Sensitivity: 2 pg/ml.   |      |
|         | Interleukin-10 – EIA - BEST   |      |
|         | Enzyme immunoassay kit for the quantitative                               |      |
| A-8774  | determination of interleukin-10 in human biological                       | 12x8 |
|         | Measurement range: 0-500 pg/ml.   |      |
|         | Sensitivity: 1 pg/ml.   |      |
|         | Erythropoietin – EIA - BEST   | 12x8 |
|         | Enzyme immunoassay kit for the quantitative                               |      |
| A-8776  | determination of erythropoietin in human biological                       |      |
|         | Measurement range: 0-200 mIU/ml.  |      |
|         | Sensitivity: 0.5 mIU/ml.  |      |